

Environmental Toxicology 2018: Toxicity response of aquatic bioindicators exposed to water polluted with glyphosate

Cristina A. Villamar

ABSTRACT

Herbicides based on glyphosate (C₃H₈NO₅P / CAS 1071-83-6) have been used (actually close to 150 million kg/year) since 1974 on crop soils to eliminate invasive plant species. Sources water can be reached by this herbicide through runoff, leaching and direct exposure of the water receiving body by aerial application. On aquatic environmental, the glyphosate has been found at concentrations up to 4.0 µg/L. Despite the glyphosate has been designed to be lethal in plants (inhibition of the shikimate pathway), the presence of surfactants from commercial formulations and metabolites formation (e.g. aminomethylphosphonic acid or AMPA) by biodegradation could change its toxicity on organisms aquatic. The aim of this work was the ecotoxicological evaluation of freshwater and marine water polluted with glyphosate on *Daphnia magna*, *Emerita analoga* and *Tisbe longicornis*. The methodology used the exposition the aquatic organisms to GLIFOPAC (480 g/L of active ingredient) at concentrations between 0.5 and 4.8 g/L of active ingredient. The acute toxicity of *D. magna* (48 h-LC50), *E. analoga* (48 h-LC50) and *T. longicornis* (96 h-LC50) were studied. Moreover, chromatographic analysis of freshwater and marine water polluted with glyphosate was evaluated. Results demonstrated that acute toxicity reported values for *D. magna*, *E. analoga* and *T. longicornis* of 27.4, 806.4 and 19.4 mg/L, respectively. Chromatographic analysis described around 45 substances of the GLIFOPAC composition such as structures from the surfactants (aliphatics chain with ester/ether group), metabolites (AMPA) and other substances (glucofuranose, glucopyranoside, galactopyranose). Preliminary assessments showed differences in the glyphosate composition within the freshwater and marine water,

which may influence the toxicity in each aquatic environment

REFERENCES

1. Altenburger R, Ait-Aissa S, Antczak P et al (2015) Future water quality monitoring – adapting tools to deal with mixtures of pollutants in water resource management. *Sci Total Environ* 512–513:540–551. <https://doi.org/10.1016/j.scitotenv.2014.12.057>
2. Anudechakul C, Vangnai AS, Ariyakanon N (2015) Removal of chlorpyrifos by water hyacinth (*Eichhornia crassipes*) and the role of a plant-associated bacterium. *Int J Phytoremediation* 17:678–685. <https://doi.org/10.1080/15226514.2014.964838>.
3. Arrhenius Å, Grönvall F, Scholze M et al (2004) Predictability of the mixture toxicity of 12 similarly acting congeneric inhibitors of photosystem II in marine periphyton and epipsammon communities. *Aquat Toxicol* 68:351–367. <https://doi.org/10.1016/j.aquatox.2004.04.002>.
4. Arts GH, Belgers JDM, Hoekzema CH, Thissen JT (2008) Sensitivity of submersed freshwater macrophytes and endpoints in laboratory toxicity tests. *Environ Pollut* 153:199–206. <https://doi.org/10.1016/j.envpol.2007.07.019>.
5. Ashauer R, Escher BI (2010) Advantages of toxicokinetic and toxicodynamic modelling in aquatic ecotoxicology and risk assessment. *J Environ Monit* 12:2056–2061. <https://doi.org/10.1039/C0EM00234H>.

Universidad Santiago de Chile, Chile;

Correspondence: Cristina A. Villamar Universidad Santiago de Chile, Chile

Received: December 07, 2021, Accepted: December 22, 2021, Published: December 28, 2021

Copyright: © 2021 Cristina A. Villamar. This work is partly presented at 18th Global Summit on Environmental Toxicology and Pharmacology September 17-18, 2018. Singapore



This open-access article is distributed under the terms of the Creative Commons Attribution Non-Commercial License (CC BY-NC) (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits reuse, distribution and reproduction of the article, provided that the original work is properly cited and the reuse is restricted to noncommercial purposes. For commercial reuse, contact reprints@pulsus.com