

# Intra-operative floppy iris syndrome for briefly browse

Burak Turgut\*

TURGUT B. Intra-operative floppy iris syndrome for briefly browse. *Oph Clin Ther.* 2018;2(1):1-2.

Intraoperative floppy iris syndrome (IFIS) has been described firstly in 2005 by Chang et al. as a triad of experienced signs associated with tamsulosin during cataract surgery including billowing/fluttering of a flaccid iris stroma in normal anterior camera irrigation current, the floppy iris tends to catch the phacoemulsification or irrigation-aspiration tip or to prolapse toward the phaco- and site-port incisions, and progressive pupil constriction [1]. IFIS has been classified as severe in which all signs of its triad are present, moderate in which billowing and pupillary constriction are observed, and mild in which only a fluttering iris is present [1-4].

IFIS is associated the use of systemic alpha-1 ( $\alpha$ -1) adrenergic receptor (AR) antagonists, especially tamsulosin, and others such as alfuzosin, doxazosin, terazosin, naftopidil, silodosin, or labetalol. IFIS increases the risk of some intraoperative and postoperative complications such as iris trauma, zonulolysis, posterior capsule rupture, vitreous loss, intraocular pressure elevation and cystoid macular edema. Risk factors for IFIS are hypertension, male sex, rivastigmine, benzodiazepines, quetiapine, 5-alpha reductase inhibitors including dutasteride and finasteride, short axial length and antipsychotic drugs [3-6]. As  $\alpha$ -1 AR antagonists are also prescribed for the female patients with lower urinary tract symptoms and arterial hypertension, IFIS can also occur in women. Recent studies reported that IFIS in patients started on tamsulosin often occurs in 9-36 months. However, IFIS may occur even with the usage of 2 days of associated drugs [3-6]. It has been reported that IFIS might occur even in the patients in which tamsulosin was ceased in two years prior to cataract surgery. The pathogenesis of IFIS has not been clearly understood. However, it is considered that the tone loss and disuse atrophy of dilator pupillary muscle due to constant  $\alpha$ -1 AR blockade; a possible additional  $\alpha$ -receptor, the 1L subtype which might mediate iris dilation affected by tamsulosin; the affection of other non-adrenergic receptors such as dopaminergic and serotonergic receptors; the drug-melanin interaction and dilator muscle atrophy due to chronic topical use of bunazosin which is a medication for the treatment of glaucoma might play role in the pathogenesis [1-8].

Following recommendations should be considered to reduce possible difficulties and complications due to IFIS and to make easy and secure, surgery in possibly IFIS cases [7-12]:

1. The surgeon should question the patient for the past and current medical history, the presence of benign prostate hyperplasia in males and the presence of the lower urinary tract symptoms and hypertension in both sexes.
2. The surgeon should predict the risk for the development of IFIS. If the amount of pupillary dilation following mydriatic drop use is less than 6 mm, the development of severe IFIS is higher.
3. Although preoperative discontinuation for 2-8 weeks prior to surgery of the drugs associated with IFIS may result in a larger pupil size at the

beginning of the operation, this cannot often prevent IFIS because the duration of the action or withdrawal period of especially tamsulosin is longer than estimated and it is an irreversible antagonist of  $\alpha$ -1 ARs. Additionally, IFIS might occur even in the patients in which tamsulosin was ceased in two years prior to cataract surgery. Thus, the discontinuation of the drugs associated with IFIS is not recommended.

4. The surgeon should consider the use of the drop combination of cyclopentolate, phenylephrine, and a nonsteroid-anti-inflammatory drug for preoperative pupillary dilation in potential IFIS cases. Additionally, the use of atropine %1 drops three times per day may be started one-three days before surgery. However, it should be careful while the use in elderly patients because of the risk of urinary retention and cardiac side effects.
5. The surgeon should consider the usage of capsular staining for easy visualization of the anterior capsulorhexis boundaries.
6. The creation of a tri-planar, self-sealing corneal tunnel incision or a slightly longer clear corneal incision, or an additional clear corneal incision, and well-constructed corneal side-port incisions may be required in potential IFIS cases.
7. The surgeon may safely use a visco-adaptive vehicle or the soft-shell technique in all IFIS cases.
8. Intraoperative intracameral phenylephrine (0.5-0.6 cc from a dilution of 1:1,000 sulfite-free) or epinephrine without preservative (0.25-0.50 cc from dilution of 1:1000 epinephrine 0.3 to 0.5 cc from dilution of 1:2500 in balanced saline solution) may be administered under the iris.
9. Addition of epinephrine to the irrigation solution may be beneficial to continue enough pupillary dilatation.
10. The use of iris retractors or iris expansion rings/instruments may be beneficial if pupillary dilation at preoperative evaluation is less than 6mm. However, sphincterotomies should be avoided.
11. The fluidics parameters such as aspiration and irrigation levels on the phaco machines (for example, the height of the bottle may be decreased approximately 10 cm) should be decreased by %10-15 percent. This lesser irrigation and aspiration is critical because it has been reported that %95 of IFIS cases were successfully managed with a combination low fluidics with a visco-adaptive device.

## REFERENCES

1. Chang DF, Campbell JR. Intraoperative floppy iris syndrome associated with tamsulosin (Flomax). *J Cataract Refract Surg.* 2005;31:664-73.
2. Cheung CMG, Awan MAR, Sandramouli S. Prevalence and clinical findings of tamsulosin-associated intraoperative floppy iris syndrome. *J Cataract Refract Surg.* 2006;32:1336-1339.
3. Davison JA. Prevention and management of intraoperative floppy iris syndrome. Dr. Farhan Zaidi (Ed.), *Cataract Surgery 2013: 5772/18122.*
4. Chang DF, Braga-Mele R, Mamalis N, et al. ASCRS White Paper:clinical review of intraoperative floppy-iris syndrome. *J Cataract Refract Surg.* 2008;34:2153-2162.

---

Department of Ophthalmology, Faculty of Medicine, Yuksek Ihtisas University, Ankara, Turkey

\*Correspondence: Turgut B, Department of Ophthalmology, Yuksek Ihtisas University, 06520, Ankara, Turkey, Tel : 90 312 2803601; Fax: +90 3122803605; E-mail: burakturgut@yiu.edu.tr

Received: February 6, 2018, Accepted: February 7, 2018, Published: February 14 2018



This open-access article is distributed under the terms of the Creative Commons Attribution Non-Commercial License (CC BY-NC) (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits reuse, distribution and reproduction of the article, provided that the original work is properly cited and the reuse is restricted to noncommercial purposes. For commercial reuse, contact [reprints@pulsus.com](mailto:reprints@pulsus.com)

5. Chatziralli IP, Peponis V, Parikakis E, et al. Risk factors for intraoperative floppy iris syndrome: a prospective study. *Eye*. 2016; 30:1039-44.
  6. Cantrell MA, Bream-Rouwenhorst HR, Watts C, et al. Pathophysiology and pre-operative evaluation of patients at risk for intraoperative floppy iris syndrome (IFIS). Dr. Farhan Zaidi (Ed.), *Cataract Surgery 2013*: 5772/19544.
  7. Flach AJ. Intraoperative floppy iris syndrome: pathophysiology, prevention, and treatment. *Trans Am Ophthalmol Soc*. 2009;107:234-9.
  8. Jaeger EA, Murchison AP, Maguire JJ, et al. Intraoperative floppy iris syndrome. In: *Willis Eye Institute 5-Minute Ophthalmology Consult*. Wolters Kluwer Lippincott Williams & Wilkins, Philadelphia 2012, 364-5.
  9. Goseki T, Shimizu K, Ishikawa H, et al. Possible mechanism of intraoperative floppy iris syndrome: a clinicopathological study. *Br J Ophthalmol*. 2008; 92:1156-8.
  10. Simaroj P, Lekhanont K, Charukamnoetkanok P. Modified surgical techniques for managing intraoperative floppy iris syndrome. *Case Rep Ophthalmol Med*. 2016; 2016: 1289834.
  11. Storr-Paulsen A, Nørregaard JC, Børme KK, et al. Intraoperative floppy iris syndrome (IFIS): a practical approach to medical and surgical considerations in cataract extractions. *Acta Ophthalmol*. 2009;87: 704-8.
  12. Sallam A, El-Defrawy H, Ross A, et al. Review and update of intraoperative floppy iris syndrome. *Expert Rev Ophthalmol*. 2011;6:469-76.
- 
-