

which is a structural counterpart of its natural substrate, yields [4,5]. A rise in drug-induced glucosylceramide in a block. There was an increase in apoptosis and a decrease in ceramide. N-(4-hydroxyphenyl)-retinamide treatment of solid tumors cell lines. A novel synthetic retinoid²³ has been discovered. Ceramide modulation is a phenomenon that occurs when the amount of ceramide in the body is GCS metabolism by these inhibitors is a potential topic for research.

Inhibitors of cyclin-dependent kinases

Cell division is a complicated process that must be properly regulated to guarantee that genetic material is faithfully divided between daughter cells. The loss of normal cell cycle control is a common occurrence in tumor cells. The activation of a family of serine/threonine kinases known as the cyclin-dependent kinases regulates the advancement of the cycle from one phase to the next (CDKs). The degradation of the CDKs' activating cyclins regulates their activity. The regulation of the transition from G1 to S phase is particularly important in chemotherapy-induced apoptosis. The Cyclin-D-CDK4 and Cyclin-E-CDK2 complexes phosphorylate and inactivate the product of the Retinoblastoma gene (Rb) during this transition [5]. The transcription factor E2F is released by phosphorylated Rb, which leads to the synthesis of gene products required for the S phase. Clinical trials have attempted to restore Rb functionality using novel small-molecule CDK inhibitors. CDK2 and CDK4 are both competitively inhibited by flavopiridol. Flavopiridol induces apoptosis in hematopoietic cell lines and has anti-leukemia and anti-lymphoma activity in xenograft models. Because of the promising results from animal models, this drug is now being tested in phase I trials at the National Cancer Institute. In addition, when applied on cell lines, flavopiridol improves the pro-apoptotic effects of various conventional medicines, and phase I trials of these combinations are under underway. UCN-01 is a protein kinase inhibitor with improved action against protein kinase C. It is an analogue of the protein kinase inhibitor staurosporine [6]. In leukemia cell lines, UCN-01 can trigger G1 arrest and activate apoptosis by hypo phosphorylating Rb. UCN-01 also inhibits the G2 checkpoint, which is activated by DNA damage, by interacting with chk1, a checkpoint kinase. UCN-01 is a protein kinase inhibitor that has been shown to be more

effective against protein kinase C. It's a protein kinase inhibitor that's similar to staurosporine. By hypo phosphorylating Rb, UCN-01 can induce G1 arrest and death in leukemia cell lines. By interacting with chk1, a checkpoint kinase, UCN-01 also blocks the G2 checkpoint, which is activated by DNA damage.

Family structures of Caspase and Bcl-2

Caspases cleave peptides after an aspartate residue and have a cysteine residue in their active site. They are made in the form of zymogens, which are precursors. The active enzyme is made up of a heterotetramer of the type 22 after the n-terminal prodomain is removed and the precursor is processed into large and small subunits simultaneously. Tetra peptides that come before the aspartate residue are recognized by caspases [7]. There are at least fourteen caspases known in mammals, which are split into three subfamilies with varying substrate preferences. The caspases are sequentially activated during apoptosis produced by diverse stimuli, constituting a caspase cascade. Anti-cancer medications and factor deprivation trigger cell death, which is inhibited by specific caspase inhibitors such Ac-DEVD-Cho and z-VAD-fmk.

Fas ligand and fas Receptor: Death factor and receptor

Cytokines play a role in cellular proliferation and differentiation. Cytokines can potentially trigger apoptosis. Tumor Necrosis Factor (TNF) is a type II membrane protein with a conserved extracellular

domain that is the prototype of a family of cytokines known as the TNF family. TNF, lymphotoxin, Fas ligand (FasL), TRAIL (also known as APO2 ligand), CD40 ligand, CD30 ligand, CD27 ligand, and OX40 ligand are all members of this family. Matrix metalloproteinase cleave most TNF family members from their membranes, resulting in soluble versions of the proteins. TNF and lymphotoxin have a jelly-roll structure and form a trimer, according to their X-ray structures, as do probably all TNF family members. Fas ligand and TNF, the apoptotic signal transduction caused by death factors has been thoroughly studied. When Fas ligand binds to Fas, the receptor becomes trimerized, which recruits pro-caspase 8 via an adaptor termed FADD/MORT1 and forms the DISC complex (Death-Inducing Signaling Complex) [8]. The DISC auto-activates pro-caspase and converts it to a mature enzyme. TNF-R1 uses the same mechanism to activate caspase with the exception that TRADD connects TNFR1 and FADD. The activated caspase can drive the apoptotic programme through two downstream mechanisms. Caspase 8 directly activates pro-caspase 3 in one route. In the other, caspase cleaves bid, a Bcl-2 family member that promotes apoptosis.

Apoptosis in the presence of other stimuli

Apoptosis is induced in a wide range of cells, including neurons and lymphocytes, by genotoxic chemicals such -irradiation and etoposide. The PI3 kinase-related ATM (Ataxia Telangiectasia, Mutated) gene is required for nerve cell death ATM recognises DNA damage as the first step in the start of apoptosis. This process is also dependent on the p53 tumor suppressor gene, suggesting that ATM phosphorylates p53 directly or indirectly. Bax-null animals are resistant to -ray-induced apoptosis in some neurons, suggesting that Bax, a pro-apoptotic member of the Bcl-2 family, is involved in this pathway as well. Bax can activate caspase by releasing cytochrome C from mitochondria. Many cells require a trophic factor to survive, and when that component is removed from the culture, apoptosis occurs. Nerve cells perish without Nerve Growth Factor (NGF), and the lack of NGF is thought to be the cause of the enormous apoptosis that occurs during neurogenesis [9]. Interleukin (IL) and colony-stimulating factors are required for the survival of many myeloid and lymphoid cells. An IL-3-dependent cell line was used to investigate the signal transduction mechanism underlying factor deprivation-induced apoptosis. Bid, another pro-apoptotic member of the Bcl-2 family, is phosphorylated and sequestered by a protein called 14-3-3 when these cells are cultured in the presence of IL-3. A protein called 14-3-3 phosphorylates and sequesters Bid, another pro-apoptotic member of the Bcl-2 family. Bid is dephosphorylated and translocated to mitochondria to cause the release of cytochrome C when the cells do not receive a signal from the IL-3 receptor. The mitochondrial pathway that leads to the release of cytochrome C appears to play a key role here as well.

Apoptotic DNA fragmentation

DNA fragmentation occurs as part of the apoptotic death process. With the identification of a specific DNase (CAD, caspase activated DNase, also known as DFF-40, DNA fragmentation factor) and its inhibitor, the molecular basis for this process was fully defined (ICAD, also called DFF-45).

CONCLUSION

Protein phosphorylation and DE phosphorylation are involved in growth and differentiation signals and cAMP and phosphatidyl inositol are tiny second-messenger chemicals. These signals are, for the most part, reversible. The apoptotic signal, on the other hand, is irreversible. The apoptotic signal activates proteases and DNase, which causes the cleavage of numerous cellular components and the morphological changes in cells and nuclei that are characteristic of apoptosis. Loss-of-function mutations in apoptosis-inducing molecules cause symptoms that show how important the apoptotic system is in

maintaining mammalian homeostasis. This system, however, is a two-edged sword. It is advantageous to remove useless, harmful, and senescent cells from our systems if they are appropriately managed. Exaggeration of this system, on the other hand, is harmful, resulting in tissue damage. It is now possible to consider a variety of apoptotic system applications to human disorders. It is now possible to consider a variety of apoptotic system applications to human disorders.

REFERENCE

1. Hickman JA. Apoptosis and chemotherapy resistance. *Eur J Cancer*. 1996;32(6):921-26.
2. Chinnaiyan AM. The apoptosome: heart and soul of the cell death machine. *Neoplasia*. 1999;1(1):5-15.
3. Lindsten T, Ross AJ, King A, et al. The combined functions of proapoptotic Bcl-2 family members bak and bax are essential for normal development of multiple tissues. *Mol cell*. 2000;6(6):1389-99.
4. Murphy KM, Streips UN, Lock RB. Bcl-2 inhibits a Fas-induced conformational change in the Bax N terminus and Bax mitochondrial translocation. *J Biol Chem*. 2000; 275(23):17225-28.
5. Yang XH, Sladek TL, Liu X, et al. Reconstitution of caspase 3 sensitizes MCF-7 breast cancer cells to doxorubicin-and etoposide-induced apoptosis. *Cancer res*. 2001;61(1):348-54.
6. Kolesnick RN, Krönke M. Regulation of ceramide production and apoptosis. *Annu rev physiol* 1998;60(1):643-65.
7. Gelmon KA, Eisenhauer EA, Harris AL, et al. Anticancer agents targeting signaling molecules and cancer cell environment: challenges for drug development. *J Natl Cancer Inst*. 1999; 91(15):1281-87.
8. Landowski TH, Qu N, Buyuksal I, et al. Mutations in the Fas antigen in patients with multiple myeloma. *J am Soc Hematol*.1997;90(11):4266-70.
9. Nagata S, Suda T. Fas and Fas ligand: lpr and gld mutations. *Immunology today*. 1995;16(1):39-43.