

# Mini Review on Two-dimensional Transthoracic Echocardiography in a Patient

Natalisa Hvizdosova\*

Hvizdosova N. Mini Review on Two-dimensional Transthoracic Echocardiography in a Patient. *Int J Anat Var.* 2023;16(1):246-247.

## ABSTRACT

**Objective:** A new CT scan showed thrombosis of this celiacomesenteric

trunk, with a lack of enhancement of the bowel wall after contrast media injection and diffuse peritoneal liquid, indicating mesenteric ischemia.

**Key Words:** Transthoracic Echocardiography; CT scan; Celiacomesenteric trunk

## INTRODUCTION

Two-dimensional transthoracic echocardiography was performed in 34(89.5) cases; Trans esophageal echocardiography, in 24(63.1); reckoned tomography, in 29(76.3); coronagraph and angiography, in 20(52.6). Preoperative shock was reported in 3(7.9) and cardiac tamponade in 18(47.4) cases. Further than half (57.9) of cases were operated on within the first 24 hours after admission. In the maturity of cases (73.7), the periphery of the aorta exceeded 4 cm. In the presence of type an aortic analysis, all cases passed surgery on cardiopulmonary bypass; its duration varied from 20 to 485 min, with a mean of 214.6 ±102.9 min. The mean aortic cross clamp time was 114.5 ±62.7min. Complete circulatory arrest was demanded in the maturity of cases (86.8), and it lasted 2 to 97 min (mean, 27.4 ±18.6 min). During cardiopulmonary bypass, body temperature was dropped to 17 - 28°C (mean, 18.9 ±1.95°C). The ideal of this study was to estimate the short-term results of surgical treatment in cases with acute aortic analysis cases and styles. A retrospective analysis of 38 cases with acute type an aortic analysis that was surgically treated at the Clinic of Cardiac, Thoracic, and Vascular Surgery, Sanitarium of Kaunas University of Medicine, from January 2004 to December 2007 was conducted. The opinion of aortic analysis was verified by employing special ways. The duration of surgery ranged from 1 to 14 hours, with a mean of 6.1 ±2.49 hours. During the early postoperative period, 12(31.6) cases failed. Postoperative bleeding was seen in 16(42.1) cases, and 6 of them failed latterly. Due to dragged bleeding, 4(10.5) cases were left with an open sternum after surgery. Resternotomy was performed in 9 cases; 3 of them failed due to multi-organ injury. During postoperative period, cardiogenic shock of colorful degrees was seen in 7(18.4) cases. Central nervous system injury passed in 9(23.7) cases. A 50-year-old man was admitted as an exigency for acute left-leg ischemia. He was a smoker and had experienced aortobifemoral bypass for atheroma 2 times before without re-implantation of the inferior mesenteric roadway (IMA). reckoned tomography (CT) angiography revealed thrombosis of the left leg of the graft as well as a superior mesenteric roadway (SMA) and a common hepato spleno gastric box arising from a celiac mesenteric box [1-2]. Successful thrombectomy was performed. Twenty-four hours later, he presented an acute abdominal pain with hemodynamic instability and elevated lactate (4 times the normal value) and transaminases (6 N). A new CT scan showed thrombosis of this celiacomesenteric trunk, with a lack of enhancement of the bowel wall after contrast media injection and diffuse peritoneal liquid, indicating mesenteric ischemia. Emergent laparotomy confirmed these radiological findings, with an ischemic liver and necrosis of the colon, the ileum and the gall-bladder. Despite an aggressive approach (thrombectomy of the SMA and Hepatic artery, total colectomy, enterectomy of 1 meter, cholecystectomy), the patient died 12 hours later [3].

The celiac box generally arises at the position of twelfth thoracic backbone and comprises the left gastric roadway, the common hepatic roadway and the splenic roadway. The superior mesenteric roadway arises 1 cm below, at the

position of L1. In embryonic development, the abdominal visceral highways arise from the primitive rearward abdominal aorta through four roots the left gastric roadway, the hepatic roadway, the splenic roadway and the superior mesenteric roadway. These roots are joined together by a longitudinal frontal anastomosis. Typically, a cleft forms in this anastomosis between the third and fourth roots therefore segregating the celiac box from the superior mesenteric roadway. The common celiacomesenteric box is a rare variation (0.5) of the anterior branches of the abdominal aorta. It's generally asymptomatic and may be discovered apropos during vascular studies or during corpse's analysis. Infrequently, it may beget symptoms including unspecific abdominal pain to real abdominal angina when associated to atherosclerosis. It was classified [4].

It is widely agreed that two visceral arteries are sufficient for digestive functions. The absence of the IMA explains the cataclysmic presentation of the thrombosis of the celiacomesenteric trunk in this case. This kind of vascular variation has to be known by surgeons and radiologists because it may change the management of abdominal aneurysms and atherosclerosis, and indicate for example the re-implantation of the IMA during aortobifemoral bypass [5]. The infection of aortic root is not common pathology; however, it is a complicated disease. Esophageal echocardiography is an informative method while diagnosing aortic root abscesses. The in hospital mortality is increased by the heart failure persisting after the operation and sepsis. The long-term survival is decreased by preoperative infective endocarditis of the prosthesis and heart failure. The mortality rate of patients older than 50 years is 3-fold higher than mortality rate of younger ones in hospital mortality rate was 14.3%. The causes of death included sustained heart failure and sepsis. All these patients were in NYHA functional class IV preoperatively; one of these patients had culture positive for *Staphylococcus aureus*. In hospital survival was 85.7% one-year postoperative survival - 80.9%, and both five-year and ten-year survivals were 76.0%. The long-term survival was negatively influenced by recurrent infective endocarditis, heart failure, and age. Death occurred in 1 patient (11.1%) of the 9 patients who at the time of surgery were younger than 50 years and 4 patients (33.3%) of the 12 who were older than 50 years at the time of operation [6-7].

Three peripherals are cannulated before the operation patient's arteries (right and left radial artery her and femoral artery) to measure BP. In addition, ponies central and peripheral intravenous fluids for infusions. The chest is opened during the procedure median sternotomy. Assessing the aortic dissection the nature of the infection, the location of its cannulation is chosen, and 4 (10.5%) patients had to cannulate their thighs artery. Two of them had to do it after the aorta delamination of the standard coronal connection during the holding operation. The third patient has this artery cannulated after diagnosing a huge ruptured ascending aneurysm of the aorta, and the fourth one during the suspension to tie aortic surgery after previous Cabrioli surgery- for whom the iliac artery is prepared before the sternotomy 24 (63.1%) cannulated aorta in the area of the arch patients, but directly into the lumen of the opened aorta (eye under control) a cannula was inserted in

Department of Anatomy, Faculty of Medicine, Pavol Jozef Safarik University in Kosice, Slovak Republic

Correspondence: Natalisa Hvizdosova, Department of Anatomy, Faculty of Medicine, Pavol Jozef Safarik University in Kosice, Slovak Republic. E-mail: natalisa.hvizdosova@upjs.sk

Received: 02-Jan-2023, Manuscript No: ijav-23-6074; Editor assigned: 05-Jan-2023, PreQC No. ijav-23-6074 (PQ); Reviewed: 19-Jan-2023, Qc No: ijav-23-6074; Revised: 24-Jan-2023, Manuscript No. ijav-23-6074; Published: 31-Jan-2023, DOI:10.37532/1308-4038.16(1).235



This open-access article is distributed under the terms of the Creative Commons Attribution Non-Commercial License (CC BY-NC) (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits reuse, distribution and reproduction of the article, provided that the original work is properly cited and the reuse is restricted to noncommercial purposes. For commercial reuse, contact [reprints@pulsus.com](mailto:reprints@pulsus.com)

10 (26.3%) patients. Myocardial protection was performed in a standard way. An increased number of 5A alleles leads to high expression of MMP-3. Thus, objective of the study was to determine whether the 5A/6A polymorphism in the promoter region of MMP-3 gene is associated with the development of dilative pathology of ascending thoracic aorta. We studied 76 patients (age ranged from 31 to 81 years; median age, 64 years) who underwent aortic reconstruction surgery due to dilative pathology of ascending thoracic aorta and a random sample of the population (n=604) aged 25-64 years, all from Lithuania [8-10]. DNA was analyzed by using real-time polymerase chain reaction to genotype polymorphism 5A/6A at a position - 1171 of the MMP3 gene promoter. The prevalence of MMP-3 genotypes was similar in the group of dilative pathology of ascending thoracic aorta and random sample of population. The frequency of 5A allele did not differ significantly between both groups and was 0.506 and 0.514, respectively. Male carriers of 5A/5A genotype were significantly younger compared with those with the 6A/6A genotype [11-13]. In conclusion, the frequency of MMP-3 promoter 5A/6A genotypes did not differ between the group of patients with dilative pathology of ascending thoracic aorta and the random sample of population, but the males with dilative pathology of ascending thoracic aorta and 5A/5A genotype required aortic reconstruction surgery at the younger age than the males carrying 6A/ 6A genotype in the MMP-3 promoter region. DNA quality and concentration were estimated by spectrophotometric analysis and by ethidium bromide stained agarose gel under ultraviolet light. The latter method was used to evaluate DNA degradation level in the samples from paraffin-embedded aortic tissue [14-15].

**ACKNOWLEDGEMENT:** None

**CONFLICTS OF INTEREST:** None.

#### REFERENCES

1. Kuo-Shyang J, Shu-Sheng L, Chiung-FC. The Role of Endoglin in Hepatocellular Carcinoma. *Int J Mol Sci.* 2021 Mar 22;22(6):3208.
2. Anri S, Masayoshi O, Shigeru H. Glomerular Neovascularization in Nondiabetic Renal Allograft Is Associated with Calcineurin Inhibitor Toxicity. *Nephron.* 2020;144 Suppl 1:37-42.
3. Mamikonyan VR, Pivin EA, Krakhmaleva DA. Mechanisms of corneal neovascularization and modern options for its suppression. *Vestn Oftalmo.* 2016;132(4):81-87.
4. Brian M, Jared PB, Laura E. Thoracic surgery milestones 2.0: Rationale and revision. *J Thorac Cardiovasc Surg.* 2020 Nov; 160(5):1399-1404.
5. Amy LH, Shari LM. Obtaining Meaningful Assessment in Thoracic Surgery Education . *Thorac Surg Clin.* 2019 Aug;29(3):239-247
6. Farid MS, Kristin W, Gilles B. The History and Evolution of Surgical Instruments in Thoracic Surgery. *Thorac Surg Clin.* 2021 Nov; 31 (4): 449- 461.
7. John C, Christian J. Commentary: Thoracic surgery residency: Not a spectator sport. *J Thorac Cardiovasc Surg.* 2020 Jun; 159(6):2345-2346.
8. Alexander AB, Xiaoying L, Clauden L. The Thoracic Surgery Residents Association: Past contributions, current efforts, and future directions. *J Thorac Cardiovasc Surg.* 2021 Sep; 162(3):917-927.e5.
9. Xin W, Bofu L. Aortic Dissection with Rare Anatomical Aortic Arch Variation Depicted by Computed Tomography Angiography. *Heart Surg Forum.* 2021; 24(2): E407-E408.
10. Foivos I, Jonathon K, Daryll B. Aberrant right subclavian artery - a rare congenital anatomical variation causing dysphagia lusoria. *Vasa.* 2021; 50(5):394-397.
11. Schizas N, Patris V, Lama N. Arc of Buhler: A lifesaving anatomic variation. A case report. *J Vasc Bras.* 2012; 37(11):9-326.
12. Penprapa SK, Brianna KR. Duplication of the inferior vena cava: evidence of a novel type IV. *Folia Med Cracov.* 2020; 28; 60(2):5-13.
13. Laurent de K, Stefano M. Variability of repairable bicuspid aortic valve phenotypes: towards an anatomical and repair-oriented classification. *Eur J Cardiothorac Surg.* 2019; 37(11):9-828.
14. Jun S, Zhang-Y, Chuan C. Postoperative neovascularization, cerebral hemodynamics, and clinical prognosis between combined and indirect bypass revascularization procedures in hemorrhagic moyamoya disease. *Clin Neurol Neurosurg.* 2021 Sep; 208:106869.
15. Qi L, Xiaojie T, Yafang D. Evaluation of Carotid Plaque Rupture and Neovascularization by Contrast-Enhanced Ultrasound Imaging: an Exploratory Study Based on Histopathology. *Transl Stroke Res.* 2021 Feb; 12(1):49-56.