

Risk factors of nocturnal enuresis

Gudisa Bereda*

Bereda G. Risk factors of nocturnal enuresis. *J Addict Clin Res* 2022;5(6):5-6.

ABSTRACT

Nocturnal enuresis is defined as the involuntary loss of urine at night, in the absence of organic (physical) disease such as urinary tract infection or detrusor over activity, at an age when a child could reasonably be expected to be dry (by consensus, at a developmental age of five yrs). Nocturnal enuresis is a significant developmental challenge for school age children and it can cause emotional and family stress and social isolation for the child. Mono symptomatic nocturnal enuresis is bedwetting occurring without any

day time incontinence or urological symptoms and it might be an explanation of a normal void occurring at an inappropriate and socially unacceptable time and place. Nocturnal enuresis resulted from the interaction of three physiological factors such as arousal disorder (defective sleep arousal); high nocturnal urine production (nocturnal polyuria); and nocturnal low bladder capacity or elevated detrusor activity, such as lack of prevention of bladder emptying during sleep, lowered bladder capacity, or bladder over activity.

Keywords: Arousal disorder; Elevated detrusor activity; Nocturnal enuresis; Nocturnal low bladder capacity; Nocturnal polyuria; Risk factors

INTRODUCTION

Nocturnal enuresis is expressed as any intermittent incontinence during a sleep in a child being at least five years old [1]. Nocturnal enuresis is a significant developmental challenge for school age children and it can cause emotional and family stress and social isolation for the child [2]. Chronic anxiety, impaired self-esteem, and delayed developmental steps such as attending camps or sleeping at a friend's house may occur as secondary challenges. The psychological and developmental injury may actually be more significant and devastating to the child than the symptom of enuresis itself [3]. Nocturnal enuresis can be classified into two groups such as mono-symptomatic nocturnal enuresis; which is bedwetting occurring without any day-time incontinence or urological symptoms and it might be an explanation of a normal void occurring at an inappropriate and socially unacceptable time and place [4]. Bedwetting correlated with day time indicators of bladder dysfunction, such as urgency or toileting frequency, is thought out poly symptomatic or non mono symptomatic nocturnal enuresis. Pediatric with non mono symptomatic nocturnal enuresis present with urge, frequency, or incontinence due to enuresis. Patients with nighttime wetting plus urge, incontinence, and frequency are considered to have poly symptomatic nocturnal enuresis [5,6]. Nocturnal enuresis is defined as the involuntary loss of urine at night, in the absence of organic (physical) disease such as urinary tract infection or detrusor over activity, at an age when a child could reasonably be expected to be dry (by consensus, at a developmental age of five years) [7]. Nocturnal enuresis is more commonly observed in pediatric and it may negatively affect the child's psychosocial development as well as interfere with the development of self-confidence and the ability to socialize [8]. Nocturnal enuresis can be separated into two types; which called as primary nocturnal enuresis: Primary nocturnal enuresis is the most common form of pediatric nighttime urinary control without a period of six consecutive months. Primary nocturnal enuresis is therefore bedwetting in a child aged five yrs or more who has never been dry for extended periods [9]. Secondary nocturnal enuresis: Pediatric with a period of six consecutive months of nighttime urinary control before incontinence. Secondary nocturnal enuresis is the onset of wetting after a continuous dry period of more than six to twelve months which can be caused by such as urinary tract infection, diabetes mellitus, spina bifida and epilepsy [10]. Family history of nocturnal enuresis, changes in antidiuretic hormone secretion rhythm, sleep disorders, retarded bladder maturity,

immature nerve development, psychology, and environment are the contributing elements for nocturnal enuresis [11].

LITERATURE REVIEW

Risk factors of nocturnal enuresis

Nocturnal enuresis resulted from the interaction of three physiological factors such as arousal disorder (defective sleep arousal); high nocturnal urine production (nocturnal polyuria); and nocturnal low bladder capacity or increased detrusor activity, such as lack of prevention of bladder emptying during sleep, lowered bladder capacity, or bladder over activity [12].

Physiological factors involved in nocturnal enuresis are discussed in turn below

Arousal disorder (defective sleep arousal): Enuresis may be more correctly analyzed as a challenge with awakening from sleep. In most pediatric with enuresis the arousal response is impaired during sleep. When urine volume exceeds bladder capacity, bedwetting occurs; although arousability is reduced, the sleep architecture itself is usually unaltered, and enuresis occurs in all sleep stages, especially during non-rapid eye movement sleep [13]. In normal pediatric, when the bladder reaches maximum capacity, there is a sudden urge for urination that does not occur correctly in enuretic pediatric. Chronic over stimulation leads to down regulation of the voiding center [14]. The risk factors for nocturnal enuresis involve long term use of disposable diaper, being male, and difficulty in awakening at night, mental stress, poverty, and family environment. With the use of disposable diaper, toilet training may be delayed [15].

High nocturnal urine production (nocturnal polyuria): Many pediatric with nocturnal enuresis have an altered diurnal rhythm of vasopressin secretion, resulting in the generation of excessive amounts of relatively dilute urine overnight. The expected bladder capacity is calculated by the formula: (age in years+1) × 30 ml, which is valid from two to twelve years. Nocturnal polyuria can be defined as greater than 130% of the expected bladder capacity for age [16]. Due to the discordance between nocturnal urine production and bladder capacity, the bladder may easily fill at night, leading to the awakening of the child for urination or in pediatric with trouble awakening, incontinence. Under normal situations, nocturnal vasopressin secretion is higher than in the daytime and leads to 50% less urine generation during the night [17]. Additionally, insufficient secretion of the

Department of Pharmacy, Negelle Health Science College, Guji, Ethiopia

Correspondence: Gudisa Bereda, Department of Pharmacy, Negelle Health Science College, Guji, Ethiopia; Tel: +251913118492; Email: gudisabareda95@gmail.com,

Received: July 04, 2022, Manuscript No. PULJACR-22-5127; **Editor assigned:** July 06, 2022, PreQC No. PULJACR-22-5127 (QC); **Reviewed:** July 20, 2022, QC No. PULJACR-22-5127; **Revised:** September 05, 2022, Manuscript No. PULJACR-22-5127 (R); **Published:** September 12, 2022, DOI: 10.37532/PULJACR.22.5(6). 5-6



This open-access article is distributed under the terms of the Creative Commons Attribution Non-Commercial License (CC BY-NC) (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits reuse, distribution and reproduction of the article, provided that the original work is properly cited and the reuse is restricted to noncommercial purposes. For commercial reuse, contact reprints@pulsus.com

antidiuretic hormone, which also leads to the production of more urine, has been seen in pediatric and the frequency of this is thought to be around two out of three pediatric [18].

DISCUSSION

Nocturnal low bladder capacity or increased detrusor activity: In pediatric with nocturnal enuresis, the emptying reflexes of the full bladder are not sufficiently prevented during sleep, resulting in involuntary voiding. The pontine micturition center of the brain stem is involved in bladder factors process [19]. Individuals with primary nocturnal enuresis have a functional bladder capacity corresponding to 70% of the expected capacity. An increase in bladder wall thickness was also resulted with ultrasonography of the identical in individuals. Pediatric with primary nocturnal enuresis, electroencephalography and cystometry records observed that bladder contractions could not be prevented in 30%-32% of enuretic pediatric [20,21].

CONCLUSION

Nocturnal enuresis is defined as the involuntary loss of urine at night, in the absence of organic (physical) disease such as urinary tract infection or detrusor over activity, at an age when a child could reasonably be expected to be dry (by consensus, at a developmental age of five years). Mono symptomatic nocturnal enuresis means bedwetting without any other lower urinary tract symptom. Nocturnal enuresis results from the interaction of three physiological factors such as arousal disorder (defective sleep arousal); high nocturnal urine production (nocturnal polyuria); and nocturnal low bladder capacity or elevated detrusor activity, such as lack of prevention of bladder emptying during sleep, lowered bladder capacity, or bladder over activity.

ACKNOWLEDGMENTS

The author would be grateful to anonymous reviewers for the comments that increase the quality of this manuscript.

COMPETING INTERESTS

The author has no financial or proprietary interest in any of material discussed in this article.

References

- Von Gontard A, Kuwertz-Bröking E. The diagnosis and treatment of enuresis and functional daytime urinary incontinence. *Dtsch Arztebl Int.* 2019;116:279-85.
- Caldwell PH, Deshpande AV, Von Gontard A, et al. Management of nocturnal enuresis. *BMJ.* 2013;47.
- Schafer SK, Niemczyk J, Von Gontard A, et al. Standard urotherapy as first line intervention for daytime incontinence: a meta analysis. *ECAP.* 2018;27:949-64.
- Dilling H, Mombour W, Schmidt MH, Schulte-Markwort E. Internationale Klassifikation psychischer Störungen: ICD-10, Kap. 2006;1991-2.
- Falkai P, Wittchen HU. American Psychiatric Association, Diagnostisches und statistisches Manual psychischer Störungen - DSM-5. Göttingen: Hogrefe;2015.
- Von Gontard A. The impact of DSM-5 and guidelines for assessment and treatment of elimination disorders. *ECAP.* 2013;22(1):61-7.
- Austin PF, Bauer S, Bower W, et al. The standardization of terminology of bladder function in children and adolescents: update report from the Standardization Committee of the International Children's Continence Society (ICCS). *Neurourol Urodyn.* 2016;35:471-81.
- Unalacak M. Enuresis nocturna prevalence and risk factors among school age children in northwest turkey. *Eur J Gen Med.* 2004;1(3): 21-5.
- Koo P. Association of obstructive sleep apnea risk factors with nocturnal enuresis in postmenopausal women. *Menopause.* 2016;23(2):175-82.
- Huang H. Prevalence and risk factors of nocturnal enuresis among children ages 5-12 years in Xi'an, China: a cross sectional study. *BMC Pediatrics.* 2020;20:305.
- Hamed A, Yousf F, Hussein MM, et al. Prevalence of nocturnal enuresis and related risk factors in school-age children in Egypt: an epidemiological study. *World J Urol.* 2017; 35(3):459-65.
- Jurkovic M, Tomaskovic I, Tomaskovic M, et al. Refugee status as a possible risk factor for childhood enuresis. *Int J Environ Res Public Health.* 2019;16(7):1293.
- Wang JJ, Wang XZ, Wang YH, et al. Effect of urinary training on urination control of infants using nappies after birth. *Chin J Appl Clin Pediatr.* 2018;33(17):1332-35.
- Mejias SG, Ramphul K. Nocturnal enuresis in children from Santo Domingo, Dominican Republic: a questionnaire study of prevalence and risk factors. *BMJ Paediatr Open.* 2018;2(1):000311.
- Tsuji S, Takewa R, Ohnuma C, et al. Nocturnal enuresis and poor sleep quality. *Pediatr Int.* 2018;60(11):1020-23.
- Savaser S, Kizilkaya Beji N, Aslan E, et al. The prevalence of diurnal urinary incontinence and enuresis and quality of life: sample of school. *Urol J.* 2018;15(4):173-79.
- Esposito M, et al. Primary nocturnal enuresis as a risk factor for sleep disorders: an observational questionnaire-based multicenter study. *Neuropsychiatr Dis Treat.* 2013;9:437-43.
- Glazener CMA, Evans JHC, Peto RE, et al. Alarm interventions for nocturnal enuresis in children. *CDSR.* 2005;2:02911.
- Arda E, et al. Primary Nocturnal Enuresis: A Review. *Nephrourol Mon.* 2016;8(4):35809.
- Doganer YC, Aydogan U, Ongel K, et al. The prevalence and sociodemographic risk factors of enuresis nocturna among elementary school age children. *J Fam Med Primary Care.* 2015;4:39-44.
- Aljefri HM, Basurreh OA, Yunus F, et al. Nocturnal enuresis among primary school children. *Saudi J Kidney Dis Transpl.* 2013;24:1233-41.