

World hepatitis day 2021 – Community awareness and screening for viral hepatitis in Accra, Ghana

Usama Kwadwo Asamoah Kusi^{1,2}, William van der Puije¹, Diana A. Asandem^{1,2,3}, Rawdat Baba-Adam^{1,2}, Hardy Agbeveve¹, Bright Asare^{1,4}, Philip Segbefia^{1,5}, Lutterodt Bentum-Ennin¹, Audrey Annan¹, Frank Osei¹, Doreen Teye-Adjei¹, Elsie Sutaya Galevo¹, Gifty Odame⁶, Gloria Ansa⁶, Linda Amoah^{1,2}, Joseph Humphrey Kofi Bonney^{2,3*}

Bonney JH. World Hepatitis Day 2021 – Community Awareness and Screening for Viral Hepatitis in Accra, Ghana. *J Emer Dis Prev Med.* 2022; 5(4):46-51.

ABSTRACT

Quantum Background: In Ghana, Hepatitis B Virus (HBV) infection remains a major public health threat as in many parts of the world. Even with an effective vaccine, there are shortfalls with low vaccine coverage among adults. To create awareness and encourage vaccination, community engagement and public-private partnerships are needed in endemic settings to help fund campaigns and offer vaccinations at no cost to under privileged people.

Objectives: An awareness and screening exercise was scheduled by University of Ghana-based Hepatitis-Malaria (HEPMAL) project team to coincide with the World Hepatitis Day (WHD) 2021. It was to engage the community in creating awareness of the menace and offer diagnostic services to ascertain prevalence levels and provide needed clinical support.

Methods: Participants from the University of Ghana community and its immediate environs were registered, taken through pre-counselling sessions where they were educated on hepatitis transmission and prevention before consenting. Eligible participants were screened for HBV markers (HBsAg, HBeAg, HBsAb, HBcAb, HbcAg) with a rapid test kit. All HBsAb-negative participants were recommended for initial vaccination at the event, whilst the subsequent shots were administered at the University Hospital Public Health Department. Hepatitis B surface Antigen-positive participants were counselled and referred for appropriate care.

Results / Outcomes: A total of 297 people, comprising of 126 (42%) males and 171 (58%) females aged between 17 years and 67 years were.

screened during the exercise. Amongst these, 246 (82.8%) showed no detectable protective antibodies against HBV and all of them agreed to and were given the first dose *HBV vaccine*. Additionally, 19 (6.4%) individuals tested positive for HBsAg and were counselled and referred to specialists from the University Hospital for further assessment and management. We found that 59 (19.9%) of our participants had previously initiated *HBV* vaccination and had taken at least one dose of the vaccine more than 6 months prior to this screening, 3 of whom tested positive for HBsAg. For the three-dose HBV vaccines deployed, a little over 20% (50/246) and a further 17% (33/196) did not return for the second and the third doses respectively, resulting in an overall 66% (163/246) of persons who completed all three vaccinations.

Conclusions / Lessons learnt: The use of different approaches including educational events and WHD activities to target groups and communities holds the key to effective general awareness-raising. A key lesson from this exercise is the need for partnerships between the health sector and corporate bodies to increase and improve awareness, care, and management of hepatitis. Additionally, home and school vaccination programmes may be adopted to enhance vaccine uptake and adherence to the vaccination schedule. We plan to extend this screening exercise to deprived and/or rural communities where HBV incidence may be higher than in urban communities. chaos and high chaos then we find the desired value inside the inter-section of both. This helps us find something in a $\aleph_3 \gg \infty$ this takes the problems around us to the next level if we are able to control a chaos then we can achieve pretty much anything.

Key Words: *Viral Hepatitis; World Hepatitis Day; HEPMAL; HBV Vaccine*

INTRODUCTION

World Hepatitis Day (WHD) is commemorated annually on 28th July with the aim of uniting the world under one theme to increase awareness and political commitment to tackle the global burden of viral hepatitis [1,2]. The theme for 2021 was “Hepatitis can’t wait”, highlighting the urgent need to eliminate hepatitis as a public health threat by 2030 amid other health concerns including the ongoing COVID-19 pandemic [3].

Hepatitis is essentially inflammation of the liver, and may be due to some medical conditions, medication, excessive alcohol use or toxins. However, the most common cause of hepatitis is a viral infection, caused mainly by five distinct hepatitis viruses (A, B, C, D and E) [4]. Viral hepatitis affects a significant number of people worldwide and results in over 1.1 million deaths annually. About 90% of the fatalities are from liver cancer and cirrhosis, which occur after decades of chronic hepatitis B virus (HBV) or hepatitis C virus (HCV) infection [5-7]. The World Health Organization (WHO) global estimates for 2019 indicate that 354 million people are living

with hepatitis B or C, with about 3 million new infections recorded annually. It is however known that less than 20% of the infected individuals’ have access to testing or treatment. Again, in Africa, a total of about 71 million people are infected with viral hepatitis, with approximately 60 million attributed to chronic HBV and 10 million to chronic HCV [8]. Estimated prevalence rates of HBV in West Africa range between 6% and 16% in the adult population, and according to reviews published in 2016 [9,10]. Ghana has a national prevalence of about 12.3%.

This high prevalence is reflective of the high burden of infection, along with a high prevalence of chronic liver disease and liver cancer (hepatocellular carcinoma). Hepatitis B and C are the leading causes of liver cancer in Ghana. The Hepatitis Foundation of Ghana reported in 2017 that there are about 4 million Ghanaians living with hepatitis B or C. Regrettably, the majority are not aware of their infection status, leaving them at risk of severe liver disease in later years [11-13]. and the potential to unknowingly spread the infections.

¹Joseph Humphrey Kofi Bonney, Department of Virology, Noguchi Memorial Institute for Medical Research, P. O. Box, LG 581 University of Ghana, Legon Accra, Ghana; ²Department of Virology, Noguchi Memorial Institute for Medical Research, University of Ghana, Legon Accra; ³Department of Biochemistry, Cell and Molecular Biology, University of Ghana, Accra Legon, University of Ghana; ⁴Department of Animal Biology and Conservation Science, University of Ghana; ⁵Department of Theoretical Medicine, Kwame Nkrumah University of Science and Technology, Kumasi; ⁶University Hospital, University of Ghana, Legon Accra.

Correspondence: Joseph Humphrey Kofi Bonney, Department of Virology, Noguchi Memorial Institute for Medical Research, P. O. Box, LG 581 University of Ghana, Legon Accra, Ghana. E-mail: Kbonney@noguchi.ug.edu.gh

Date of Submission: 20-Sep-2022, Manuscript No. puljedpm-22-5369; Editor assigned: 22-Sep-2022, Pre QC No. puljedpm-22-5369 (PQ); Reviewed: 25-Sep-22, Qc No. puljedpm-22-5369 (Q); Revised: 27-Sep-2022, Manuscript No. puljedpm-22-5369(R); Published: 29-Sep-2022, DOI No: 10.37532/puljedpm.2022 .5(4); 46-51.



This open-access article is distributed under the terms of the Creative Commons Attribution Non-Commercial License (CC BY-NC) (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits reuse, distribution and reproduction of the article, provided that the original work is properly cited and the reuse is restricted to noncommercial purposes. For commercial reuse, contact reprints@pulsus.com

Notwithstanding the high global burden of viral hepatitis and steady efforts being made to improve treatment and care, many people infected remain unaware of their status until they advance in the disease stage. It is established that most low-income countries have only about 5% of infected persons knowing their status. This makes others vulnerable as their ignorance makes them prone to transmit the infection to others around them [14,15]. This brings up the necessity for more subsidized HBV and HCV screening with better policies and testing standards at health facilities [15,16] and for exploring the use of other hepatitis awareness-creating approaches that target groups and communities.

In Ghana, HBV is regarded as one of the infections of public health significance and hence warrants matching attention¹⁰. A safe and effective vaccine that offers 98% to 100% protection against hepatitis B is available. Vaccination programmes were first introduced and have been available since 1992, 11 years after the first vaccine was approved for use in the USA. They come in the form of monovalent formulations for birth doses or for vaccination of at-risk adult populations like health workers, though not for free [14,17]. The standard regimen is 3 doses; the second given a month after the first dose, and the third given 6 months after the first dose [18,19]. Vaccinations for HBV which is a key recommendation to control the infection has proven to be very efficient in conferring immunity [20,21]. Hitherto HBV and HCV infections can be prevented, controlled and treated - while HBV can be treated, HCV is even curable. This notwithstanding, over 90% of people living with hepatitis B and C in Africa still do not have access to the needed care [8,22,23].

A planned awareness and screening exercise was scheduled to coincide with the World Hepatitis Day (WHD) by a University of Ghana-based Hepatitis-Malaria (HEPMAL) project which is funded by the European and Developing Countries Clinical Trials Partnership (EDCTP) and has an objective to investigate the clinical and immuno-pathological consequences of chronic HBV and *Plasmodium falciparum* co-infections [24,25]. The research group collaborated with recognized bodies and institutions including the Ghana Health Service, Hepatitis Society of Ghana (HEPSoG), Ghana Association for the Study of Liver and Digestive Diseases (GASLIDD), the Hepatitis Foundation of Ghana as well as funding partners such as AngloGold Ashanti to set up various activities to commemorate WHD 2021. Among the prearranged activities was a medical outreach programme which entails education on viral hepatitis, free screening for HBV, as well as free HBV vaccination for screened and eligible individuals.

Materials and method

The University of Ghana campus has a student population of about 38,000 when school is in session and close to 5000 members of the community belonging to the staff, their families and others plying their trade within the main campus. Within this community is a sprawling market located centrally around a collection of banks in an area called the 'banking square'. It is in this area that we mounted three 10 x 10 compact 40-inch canvas canopy tents for our walk-in service. We set up to provide to interested members of the community screening and vaccinating for HBV at no cost to them. They were first taken through a registration process where details including name, contact numbers and previous vaccination history were taken. The brand of the vaccines we administered was the Engerix-B (Hep-B) - noninfectious recombinant DNA vaccine containing hepatitis B surface antigen (HBsAg). It is produced by Glaxo Smith Kline (GSK), UK, from genetically engineered yeast (*Saccharomyces cerevisiae*).

Clinicians and nurses from the University of Ghana Hospital (UGH), the University of Ghana Medical Centre (UGMC), the Korle-Bu Teaching hospital (KBTH) and the Greater Accra Regional Hospital were in attendance to provide education on viral hepatitis, pre- and post-vaccination counselling services, and to administer the vaccines to eligible individuals. Registered individuals were given a brief background of the purpose of the event and educated on viral hepatitis including the mode of transmission, types and stages of the disease condition including complications resulting from the infection. Eligible volunteers who consented were then taken through a pre-test counselling session before 2 ml of venous blood was collected into a sterile EDTA vacutainer tube with a single use butterfly needle blood collection kit.

Haemoglobin levels were determined using a portable haemoglobin meter (URIT-12 Haemoglobin Meter, Accurex Biomedical Pvt. Ltd, Mumbai, India) with the recommended test strips. A Ministry of Health

(MoH) / Ghana Health Service (GHS) recommended HBV rapid test strip, 'The Advanced Quality One-Step multi-HBV test kit' (InTec Products Inc, Xiamen, PRC) was used to determine the HBV status of the participants. The test panel was used to determine the HBsAg, HBsAb, HBeAg, HBcAg and HBeAb status of the participants. Those that tested negative for both HBsAg and HBsAb all accepted and were vaccinated, while those that tested negative for HBsAg but positive for HBsAb were not vaccinated since they already had antibodies against the virus. A rigorous follow-up schedule of making telephone calls a week before and on-the-day reminders to all vaccinees for the second and the third doses was prearranged. Those that tested positive for only HBsAg were taken through a post counselling session and were booked for hospital appointments at the UGH.

RESULTS


Chaos we In all, 297 participants between the ages of 17 and 67 years attended the event, 126 (42%) of whom were males and 171 (58%) were females (Table 1). Of this number, 283 were screened, amongst which 19 (6.4 %) tested positive for HBsAg, all of whom were also positive for HBeAg and HBcAb. Additionally, 17 (5.7 %) tested positive for HBsAb, out of which one also tested positive for HBcAb. From all the participants screened, only 1 person tested positive for HBeAb and negative for all other antigens and antibodies. Twenty-one (7.1 %) of the participants tested positive for HBcAb, two of whom were negative for HBsAg, HBeAg and HBsAb; only one was positive for HBsAb.

A total number of 59 (19.9 %) of the eligible volunteers had previously taken the HBV vaccination. Three of the 59 tested positive for HBV surface antigen. The first of the 3 doses of the HBV vaccine were successfully administered to 246 of participants who tested negative for all HBV antigens and antibodies. By the end August 2021, which was the due date to complete vaccination for the second doses, 196 (79.7 %) of the 246 who took the first dose had taken their second dose. This meant 50 (20.3 %) individuals were lost to follow-up. The final dose which was administered six months (January 2022) after the first dose registered 163 vaccinees out of the 196 eligible recipients with 33 (16.8 %) individuals lost to follow-up. Vaccine success rate overall was therefore 66%.

Table 1: Results summary

Category	Results				
	Male	Female	Total		
Number of participants	Gender	126	171		
	Mean Age (Range)	31.5 (19-67)	28.4 (17-64)	297	
Vaccination history	Vaccinated*	18	41	59	
	Not vaccinated	107	129	236	
	Unspecified #	1	1	2	
HBV Profile	HBsAg	Positive	14	5	19
		Negative	108	156	264
	HBsAb	Positive	6	11	17
		Negative	116	150	266
	HBeAb	Positive	0	1	1
		Negative	122	160	282
	HBeAg	Positive	14	5	19
		Negative	108	156	264
	HBcAb	Positive	15	6	21
		Negative	107	155	262

Shows people who have previously initiated vaccination, i.e., taken at least 1 dose of the vaccine prior to this screening exercise # Did not say whether or not they have received HBV vaccination prior to this exercise



DISCUSSION

Now what The WHO's Global Health Sector Strategy (GHSS) on viral hepatitis hopes to achieve elimination of viral hepatitis by 2030. This means reducing the annual disease incidence and mortality by 90% and 65% respectively using 2015 data as baseline¹⁰. Diagnosis and awareness of infection is part of the first essential steps towards achieving this goal. The HEPMAL Project, a University of Ghana-based EDCTP2-funded study of the immunopathological effects of HBV and malaria co-infections, sought to address and support this drive with an HBV educational, screening and vaccination exercise on the University of Ghana's main campus. Our screening and vaccination exercise examined the awareness rate among the volunteers and the prevalence of HBV within the University community. We established from the eligible volunteers a suboptimal rate in awareness through our counselling sessions and an active case prevalence of 6.4%. This indicates ongoing active viral replication within these members of the community who hitherto were not aware and hence have a high propensity to transmit the virus to vulnerable individuals. This prevalence rate is comparable to the estimated 8.36% national prevalence for adults in a systematic review and meta-analysis of HBV infections between 2015 and 2019 in Ghana [26]. The same study reported prevalence rates of 0.55% and 14.30% among pre-school children and adolescents, respectively [26].

In Ghana, HBV is the most dominant cause of viral hepatitis and requires immediate attention, especially through rapid implementation of the birth dose HBV vaccines, active case search for timely intervention, and vaccination of uninfected adults. We used *Engerix-B* HBV vaccines for our exercise. This is a 3-dose series with 2 doses separated by ≥ 4 weeks, and a 3rd dose 4 months to 6 months after the 2nd dose. An individual lost to follow-up for not completing the series, for example missing out on the second and/or the third dose, the series does not need to be restarted. Though we had a pre-counseling session with the needed information on the vaccines including the doses-scheduling for all vaccinees, our vaccination coverage for all negative anti-HBV antibody volunteers was not completed. We lost 50 (20.3%) and 33 (16.8%) persons to follow-up for the second and third dose series respectively. Through the rigorous call up and reminder schedules we set up for vaccinees prior to their return for the second and the third dose series, we had various reasons including reallocation and simply making time to come over for the dose. Overall, we achieved a full vaccination success rate of about 66%, largely due to the follow-up strategy that was implemented.

Data gathered showed that 19% of participants had previously had HBV vaccines but indicated they did not complete the three-dose regimen. They took at least one dose of the vaccine more than 6 months prior to this screening. Three out of this number had since become infected with the virus, testing positive for HBsAg alone. These are non-responders as they have not developed protective antibodies, and this may be attributed to their inability to complete the full series of the vaccines. Eligible volunteers who passed for vaccination during our event were advised of the importance of completing the series of the vaccine regimen. Even with the counseling, 20% of the of the number that took the first dose of the vaccine did not show up for the second dose. This may be attributed to various reasons including genuine inability to be present due to residency but may not be hesitancy for the vaccines. Control programmes may need to explore home or school vaccination strategies as is implemented for the polio vaccine in developing countries to minimize losses to follow-up.

Another interesting observation is the finding that a higher number of males (60) were lost to follow up than females (23), even though more males signed up for vaccination at the beginning. This may be important for informing gender-oriented follow-up strategies that will improve upon full vaccine uptake.

The Ghana Health Service (GHS) is on the verge of implementing HBV birth dose in health facilities nationwide, which can be complemented with active surveillance activities as opposed to the current passive surveillance method employed to accelerate HBV elimination. This will identify infected people who may be living apparently healthy lives for timely intervention to increase their life expectancy and curb community transmission. Additionally, we recommend private sector partnerships to assist in funding such laudable endeavors and extending them beyond WHD commemoration events. Our findings also emphasize the need to increase awareness creation for viral hepatitis in Ghanaian communities even among the literate to encourage voluntary vaccinations among teenagers and adults who were born before the HBV birth dose was implemented. We encourage more of such programmes and hope to partner with other corporate institutions to extend this to rural communities as well.

The use of screening strategies in communities to target groups is expected to be vital to the goals for general awareness-raising. Assessment of the effectiveness of this many-sided approach is necessary to reduce the undiagnosed population and improve the link between testing and care for viral hepatitis.

Acknowledgement

We acknowledge the contribution of Dr Atsu Godwin Seake-Kwawu and the Hepatitis Desk of the Ghana Health Service, members of the Hepatitis Society of Ghana (HEPSoG), Ghana Association for the Study of Liver and Digestive Diseases (GASLIDD) and the Hepatitis Foundation of Ghana. We also thank nursing officers Patience Attipoe, Anita Tay, Enyonam Baah-Danquah, Franklina Aboagye and Sammy Gyedu for their help.

Funders

We were immensely supported by AngloGold Ashanti Limited, a global gold mining company. We also had funding support from the HEPMAL project – an European Commission EDCTP2-funded project based at the University of Ghana, the West African Centre for Cell Biology of Infectious Pathogens (WACCBIP), Inqaba Biotec West Africa Limited (IBWA) and Kingdom Books and Stationary (KBTS) Limited. We also received first and second dose series vaccines free of charge from Crescent Chemist Co. Limited, agents of Biotest Pharma of Germany and we are indeed grateful to them.

REFERENCES

- Laskar MS. Different health related days in a calendar year observing worldwide. *Med*. 2018 ;5(1):1-4.
- Lavanchy D. Viral hepatitis: global goals for vaccination. *J Clin Virol*. 2012 ;55(4): 296-302.
- Lakra S. World hepatitis day: Hepatitis can't wait 28 July.
- Hooper J, Martin M. Overview of hepatitis B and C management. *US Pharma*. 2009;34(12): 42-44.
- Ajuwon BI, Yujuico I, Roper K, et al. Hepatitis B virus infection in Nigeria: a systematic review and meta-analysis of data published between 2010 and 2019. *BMC infect dis*. 2021;21(1): 1-5.

6. World Health Organization. Global progress report on HIV, viral hepatitis and sexually transmitted infections, 2021: accountability for the global health sector strategies 2016–2021: actions for impact: web annex 2: data methods.
7. World Health Organization. World Hepatitis Day 2019. Retrieved in February 2022, <https://www.who.int/campaigns/world-hepatitis-day/2019>.
8. Ismail F, Haq S, El-Garawani I, et al. Hepatitis C Virus Infection in Eastern Libya: Efforts Needed to Improve HCV Testing and Linkage to Care in the Resource-Limited Setting. *Trop Med Infect Dis.* 2022 ;7(2): 14.
9. Coste M, De Sèze M, Diallo A, et al. Burden and impacts of chronic hepatitis B infection in rural Senegal: study protocol of a cross-sectional survey in the area of Niakhar (AmBASS ANRS 12356). *BMJ open.* 2019 ; 9(7): 302-311.
10. Ofori-Asenso R, Agyeman AA. Hepatitis B in Ghana: a systematic review & meta-analysis of prevalence studies (1995-2015). *BMC infect dis.* 2016;16(1): 1-5.
11. Jandura S, MacLachlan, J.H. and B.C.J.C.S.H.p.i.m. Cowie. Hepatitis B virus epidemiology. 2015.5(5): 214-215.
12. Wang T. Model of life expectancy of chronic hepatitis B carriers in an endemic region. *J epide.* 2009 ;19(6): 311-318.
13. Yamoah P, Mensah KB, Attakorah J, et al. Adverse events following immunization associated with coronavirus disease 2019 (COVID-19) vaccines: A descriptive analysis from VigiAccess. *Hum. Vaccines Immunother.* 2022; 2109-2365.
14. Osei E, Niyilapah J, Kofi Amenuvege G. Hepatitis B knowledge, testing, and vaccination history among undergraduate public health students in Ghana. *Bio res intern.* 2019; 2019:1-3.
15. Helegbe GK, Tanko F, Aryee PA, et al. High hepatitis b seroprevalence, low knowledge, and poor attitude towards hepatitis b virus infection among market women in Bolgatanga Metropolis in the Upper East Region of Ghana. *J Trop Med.* 2020 May 27; 2020.
16. Coffie PA, Egger M, Vinikoor MJ, et al. Trends in hepatitis B virus testing practices and management in HIV clinics across sub-Saharan Africa. *BMC infect dis.* 2017;17(1):141-148. [
17. Moticka EJ. A historical perspective on evidence-based immunology. *Newnes;* 2015; 2(5):5-9.
18. Kumah A, Tormeti E, Dzando G, et al. Knowledge, Attitude and Practices towards Hepatitis B Infection and Vaccination among Public Health Students in Ghana. *Open J Prev Med.* 2005;11(1): 43-53.
19. Hou, J., Z. Liu, and F.J.I.j.o.m.s. Gu, Epidemiology and prevention of hepatitis B virus infection. *Int J Med Sci.* 2005; 2(1): 50.
20. Meireles LC, Marinho RT, Van Damme P. Three decades of hepatitis B control with vaccination. *World J hepatol.* 2015;7(18): 21-27.
21. World Health Organization. Combating hepatitis B and C to reach elimination by 2030: advocacy brief. World Health Organ; 2016.
22. Padmasola V, Ahmed MZ, Ahmed O, et al. Epidemic of COVID-19 in China and associated psychological problems. *Asian j psychiatry.* 2020; 51:102092.
23. Cox AL, El-Sayed MH, Kao JH, et al. Progress towards elimination goals for viral hepatitis. *Nat Rev Gastroenterol Hepatol.* 2020;17(9):533-42.
24. Clinical and Immunopathological Consequences of Chronic HBV and Plasmodium falciparum Co-infections. *HEPMAL.* 2020; 2(2).
25. EDCTP Senior Fellowships. EDCTP. 2018. 2020.
26. Abesig J, Chen Y, Wang H, et al. Prevalence of viral hepatitis B in Ghana between 2015 and 2019: A systematic review and meta-analysis. *PloS one.* 2020;15(6):e0234348.